Name, Date, Hr/Per 3. When the concentration of solute inside & outside a cell is the same, the cell has reached A. maximum concentration **Cellular Transport Worksheet** C. osmotic pressure Answer the following questions using your notes and your textbook. 4. The diffusion of water across a selectively permeable membrane is called **OSMOSIS** - Write the correct type of solution underneath (isotonic, hypertonic, or hypotonic) A. active transport C. osmosis 5. Energy for active transport comes from a cell's A. Golgi complex C. mitochondria 3. 6. transport requires energy from ATP to move substances across membranes. A. Passive tonic there is a GREATER concentration of solute molecules OUTSIDE the cell than inside. 7. All of the following are kinds of passive transport EXCEPT A. Diffusion 5. tonic B. Osmosis there is a LOWER concentration of solute molecules OUTSIDE the cell than inside. 8. When molecules move DOWN the concentration gradient it means they're moving from tonic 6. A. an area of low concentration to an area of higher concentration there is the SAME concentration of solute molecules outside the cell as inside. B. an area of high concentration to an area of lower concentration Fill-Ins - Complete the transport terms. Some of the letters have been filled in! 7. The SWELLING AND BURSTING of animal cells when water enters I. Active transport requires <u>E</u> ______ to move molecules across membranes. happens when a cell is placed in a tonic solution. 2. <u>A</u> _____ is the molecule that provides the energy for active transport. 3. D_____ moves oxygen and carbon dioxide molecules from a high 8. What organelle [that plants have that animals don't] keeps plant cells from Cells swell and burst bursting in this condition? concentration to a low concentration across membranes. 4. The cell organelles that burns glucose and provides ATP for active transport are the 9. The SHRINKING of plant cells when water leaves so the cell membrane pulls <u>_M_</u>_____ 5. Water moves across membranes by <u>O</u>_____. away from the cell wall or shrinking of animal cells happens when a plant cell is placed into a tonic solution. Cells shrink 6. A small membrane sac used to transport substances during exocytosis & endocytosis and shrivel 10. Cells stay the same size when placed in an tonic solution because the amount of water leaving the cell is the same and the amount of water entering. concentration of molecules outside the cell is the same as inside. 9. A solution in which there is a HIGHER concentration of molecules OUTSIDE the cell than **MULTIPLE CHOICE:** Circle and/or fill-in the answer(s) that best completes the sentence. inside = H I. The substance that dissolves to make a solution is called the 10. A CONCENTRATION <u>G</u>______ forms whenever there is a difference in A. diffuser B. solvent C. solute D. concentrate concentration between one place and another. 11. A solution in which the concentration of molecules outside the cell is LOWER than inside 2. During diffusion molecules tend to move C. down / with the concentration gradient A. up / against the concentration gradient 12. When molecules move from high to low along a concentration gradient we say they are B. from an area of lower concentration to an D. in a direction that doesn't depend on moving "<u>D</u>____" the gradient. area of higher concentration concentration

B. homeostasis

D. phagocytosis

B. nucleus

B. facilitated diffusion


D. ion channels

B. Active

D. lysosomes

D. dynamic equilibrium

B. facilitated diffusion




2. If the solute (dots) in this diagram is unable to pass through the dividing membrane, what will happen?A. the water level will rise on the right side of the tube

- B. the water level will rise on the left side of the tube
- C. the water level will stay equal on the two sides

Match the description with the solution type:

A. Isotonic B. Hypertonic C. Hypotonic	I solution with a lower solute concentration (more water)
	2 solution in which the solute concentration is the same
	3 condition plant cells require
	4 condition that animal cells require
	5 red blood cell bursts (cytolysis)
	6 plant cells shrink (plasmolysis)
	7 solution with a higher solute concentration (less water)
	8 solution with a high water concentration

Label the tonicity for each solution (isotonic, hypotonic, or hypertonic): Pay close attention to the arrows!!!



Examine the pictures on the bottom of the left side of this page. What [if anything] is different about the plant and animal cells in each of these states?

<u>State</u>	Animal Cell	Plant Cell
Hypertonic		
lsotonic		
Hypotonic		

Matching - Match each term to its definition.

a. energy	I. Transport protein that provides a tubelike opening in the plasma
b. facilitated diffusion	membrane through which particles can diffuse 2. Is used during active transport but not passive transport
c. endocytosis	3. Process by which a cell takes in material by forming a vacuole
d. passive transport	around it
e. active transport	4. Particle movement from an area of higher concentration to an area of lower concentration
f. exocytosis	 5. Process by which a cell expels wastes from a vacuole 6. A form of passive transport that uses transport proteins
g. protein ion pump	7. Particle movement from an area of lower concentration to an
h. channel protein	area of higher concentration 8. Transport protein that changes shape when a particle binds
	with it
Chant Annuan	

Short Answer –

I. Name two factors that affect the rate of diffusion.